

INSTRUCTIONS FOR THE USE OF ALLERGENIC EXTRACTS ALLERCEPT® THERAPY DROPS

Therapy

ALLERCEPT® Therapy Drops are indicated as an aid in alleviating symptoms associated with allergic disease in the dog, cat and horse. Other factors in the management of the allergic animal include environmental control and elimination of the offending allergens when possible.

Three Bottle Sequence

The drops come in 3 formulations (A, B and C). Your first prescription will contain two bottles, one each labeled "A" and "B". Begin treatment with bottle "A". Once you have completed this bottle, begin bottle "B". Your second prescription order as well as subsequent refills will contain two bottles labeled "C". After you have used bottle "B", begin bottle "C" and continue with this for the duration of treatment.

Priming the Dispenser

To begin using the dispenser for any new bottle, IT IS IMPORTANT THAT YOU FIRST PRIME THE PUMP by removing the colored plastic safety clip and depressing the pump several times firmly and quickly until a dose is released.

Warning: Keep the safety clip away from small children and pets as it may pose a choking hazard.

Store bottles at room temperature in an upright position. If your bottle is unused for more than a day or two, you may need to re-prime it to ensure doses are properly released.

Dispensing the Drops

Rest the dispensing arm of the pump on the lower teeth, anywhere near the front or side of the mouth, with the tip of the arm over the edge of the teeth (Figure 1). To administer, fully depress the pump firmly and quickly so that the dose is released into the mouth, preferably under the tongue. Repeat a second depression of the pump for a total of two pumps per dose. It is important to keep the bottle in an upright position while depressing the pump. It may be helpful to gently hold your pet's mouth closed while administering the dose (Figure 2).

Do not mix the drops with food or on a treat. The drops are highly palatable for most pets. Avoid giving food or water for 10 minutes after dosing to allow the drops to remain in the mouth as long as possible.



Figure 1



Figure 2

Daily Dosing

Administer the 2-pump dose twice per day, for example, morning and evening. The doses do not have to be given at the same time every day. If you miss a day, do not try to "catch up" by giving additional doses; just continue treatment the next day. Each bottle will last approximately 75 days. Order refill bottles through your veterinarian approximately 2–3 weeks before needed.

Warning: In the unlikely event that your pet experiences a reaction, contact your veterinarian. Do not resume drops until you have discussed your pet's reaction with your veterinarian.

Manufactured by

Diamond Animal Health, Inc.
Des Moines, IA 50327 U.S.A.
U.S. Veterinary License No. 213

Distributed by

Heska Corporation
Loveland, CO 80538 U.S.A.
1-800-464-3752

